fresh tomatoes/capsicums

Reduces transplant shock

8

- Increases root number and shoot vigor
- Improves resistance to nematode infestation
- Increases fruit size and number, and total yield
- Produces higher early harvest yields
- Increases shelf-life of fruit by up to 1 week



KELPAK

O°

| - | the | |
|---|-----|---|
| | - | |
| | _ | / |

| Kelpak on tomatoes | | | | | |
|--------------------|--------------------|---------------------------|-----------------------------|-------------------------------------|--|
| COUNTRY | ТҮРЕ | RATE (pt/Ac) | APPLICATION | AVE. YIELD INCREASE ¹ | |
| California | Processing | 1.0% 2 - 2.5 pt/Ac x 2 | Dip/drench Foliar sprays | 10% | |
| Chile | Greenhouse | 1.0% 0.5% x 3 | Dip/drench Foliar sprays | 23% | |
| Hungary | Field | 1.0% 2 pt/Ac x 2 | Dip Foliar sprays | 21%* | |
| Philippines | Field | 1.0% 2 pt/Ac x 3 | Dip Foliar sprays | 31% | |
| Poland | Field | 2 pt/Ac x 3 | Foliar sprays | 7% | |
| South Africa | Greenhouse & Field | 1.0% 2 pt/Ac x 3 - 5 | Dip/drench Foliar sprays | 23% | |
| Spain | Greenhouse | 7 pt/Ac x 1 | Drip after plant | 70%* | |

¹ Yield increase above standard grower program

* Yield of first 3 pickings

| Shelf-life of tomatoes treated with Kelpak | | |
|--|------------------------|--|
| APPLICATION | IMPROVED Shelf-Life | |
| 3 x foliar | + 5 days | |
| 5 x foliar | + 7 days | |
| Soil drench | + 6 days | |
| Dip + 3 x foliar | +10 days | |

| Y | 3 |
|---|---|
| | |

| Kelpak on capsicum crops | | | | |
|--------------------------|------------|-------------|-------------------------------------|--|
| COUNTRY | ТҮРЕ | VARIETY | AVE. YIELD INCREASE ¹ | |
| South Africa | Greenhouse | Rodeo | 26%* | |
| South Africa | Greenhouse | King Arthur | 40%* | |
| South Africa | Field | Kent Yellow | 14% | |
| California | Field | Excalibur | 14%** | |
| Peru | Field | Resistant | 13% | |

¹ Yield increase above standard grower program

* Marketable fruit from fruit picked in a single harvest

** Increase in Jumbo size fruit



RECOMMENDED APPLICATION RATE

Dip the roots of seedlings (or seedling tray) in 2 pts Kelpak per 25 gal water before transplanting into the field or greenhouse

Follow up with a 2 pt/Ac Kelpak foliar spray 14 days later and repeat the foliar spray once or twice at 14 day intervals

Spray direct seeded crops at 3 to 4-leaf stage and repeat once or twice at 14 day intervals

Alternatively to seedling dip at plant-out, Kelpak may be applied at 6-8 pt/Ac as a pulse through drippers after transplanting. Rinse lines after pulse