
cherries

- Improves pollen germination
- Improves pollen tube growth
- Increases fruit set and fruit retention
- Reduces post-harvest fruit split
- Increases fruit size and weight

- El alas

- Improves marketable yield
- Improves fruit quality













Effect of Kelpak on cherry yield					
LOCATION OF STUDY	APPLICATION RATE	VARIETY	YIELD (ton/ha) CONTROL KELPAK		INCREASE (%)
California	3-5 sprays at 300 ml/100 L	Bing	10.8	14.8	37
France	3 sprays at 300 ml/100 L	Lapin	6.5	8.5	31
Chile	3-5 sprays at 300 ml/100 L	Bing	12.7	16.5	30
Chile	3 sprays at 300 ml/100 L	Lapin	26.4	28.3	7

Kelpak on cherry pollen growth					
TREATMENTS	POLLEN GERMINATION	POLLEN TUBE LENGTH			
	%	μm			
Control	47.0 ± 1.6 b	71.3 ± 9.2 b			
Kelpak	64.0 ± 3.5 a	128.6 ± 9.2 a			





RECOMMENDED APPLICATION RATE

Spray 3 to 5 times at a rate of 300 ml/100 L starting at 30% bloom with 10 day intervals

Kelpak can be applied in tank mixes with other agrochemicals. Keep the pH of the spray solution below 7 for optimum results Kelpak is manufactured using the unique cold Cellburst extraction process







KELPAK